

Going Big in Health: Effect of a Large-Scale Preventive Health Policy^{*}

Arturo Aguilar[†]

Ricardo Gomez-Carrera[‡]

Adrian Rubli[§]

May 7, 2025

This paper studies how a conditional cash transfer program with an embedded healthcare component might induce relevant tradeoffs. By improving a previously employed identification that exploits a sharp discontinuity in PROGRESA's initial locality-level rollout, we estimate a sizable 12% increase in outpatient visits at public clinics, mainly driven by children and adults aged 20-49. This translates into improvements in reproductive healthcare and screenings for chronic diseases. While the attendance of non-targeted elderly members is not affected, the program does seem to impose costs by increasing congestion and reducing self-reported quality. This suggests that the benefits of this policy lever may carry negative unintended effects to the non-beneficiary population.

JEL Codes: I18, I15

Key words: preventive health, conditional cash transfers, health utilization, congestion.

^{*}We thank Andrew Foster, Elaine Liu, Craig McIntosh and the seminar participants at LACEA 2022, MWIEDC 2023, iHEA Congress 2023, WEAI 2023, and ITAM-ND Miniconference 2023 for their helpful comments. We acknowledge support from the Asociación Mexicana de Cultura. Part of this research was conducted while Ricardo Gomez-Carrera worked at Banco de México; the views and conclusions presented in this paper do not necessarily reflect those of Banco de México or its Board of Governors. All errors are our own.

[†]Centro de Investigación Económica and Department of Economics, ITAM. Email: arturo.aguilar@itam.mx.

[‡]World Inequality Lab. Email: ricardo.gomez-carrera@psemail.eu.

[§]Department of Business Administration, ITAM. Email: adrian.rubli@itam.mx.

1 Introduction

Across a variety of contexts, the existing literature has estimated high returns to preventive health. Despite this, uptake is typically low and highly sensitive to prices, even in situations where it is considerably subsidized (Dupas and Miguel, 2017). One potential mechanism for encouraging its adoption is to embed it as a condition for other desirable (and potentially more salient) benefits. Interventions such as school vaccine mandates (Lawler, 2017; Abrevaya and Mulligan, 2011), workplace or employer-based wellness programs (Cawley and Price, 2013; Jones et al., 2019) and conditional cash transfer programs (Levy and Ohls, 2010) are examples of these policy levers.

In this paper, we take advantage of the national expansion of the PROGRESA conditional cash transfer (CCT) program in Mexico from 2000 to 2003 to estimate its impact on health behaviors, health outcomes, and possible negative spillovers to the non-targeted population.¹ The program required all family members of a recipient household to attend preventive healthcare visits regularly.² During its initial roll-out, the program was expanded in disadvantaged rural locations, reaching over 4.2 million households in 70,436 localities by 2003, which amounts to approximately 30% of the population living in the areas served by the program (Hernández Licona et al., 2019). At its peak, the program reached 6.5 million households (equivalent to 18% of total households in Mexico), meaning that the increase in potential demand for health services created by the program is sizeable.

The questions we seek to answer in this paper are threefold. First, using administrative information, we verify that the program indeed led to a significant increase in the demand for health services at public clinics and explore the demographic composition of said increase. Second, we study which measures of preventive healthcare and behaviors changed due to the program. Finally, we analyze if the sudden increase in demand — which was unaccompanied by an increase in the supply of public clinics — led to congestion of health services, potentially increasing the cost of attending for non-beneficiaries of the program and crowding-out their healthcare utilization.³

¹PROGRESA is a well-known and widely researched CCT program that was implemented in Mexico starting in 1997 and expanded nationally thereafter.

²This requirement varied in frequency depending on the age, gender, and condition of each member. For example, pregnant and breastfeeding women were required to attend more frequently. See online appendix Table A1 for details.

³The implementation of the program acknowledged the need for strengthening the supply of health services to meet the additional demand (Secretaría de Desarrollo Social, 2000). However, the government only devoted resources to improving staffing and medical supply needs at the existing clinics, without increasing the actual supply of clinic services (Skoufias, 2001).

To identify the causal impact of the program, we employ a regression discontinuity analysis. We mostly follow the strategy in [Alix-Garcia et al. \(2013\)](#), which consists of exploiting an administrative rule that defined how the program would be rolled out at the locality level. However, instead of relying on a single cutoff rule, as previously done, we use detailed administrative records and a data-driven procedure to identify region-specific cutoffs during our sample period. To be included in the program, localities were required to have access to public clinics and schools (to be able to enforce the conditionality) and had to be poor, as defined by a pre-specified government-measured index. This index and the corresponding regional cutoff levels form the basis for the discontinuity during the first years of PROGRESA’s roll-out. Our first stage yields a 35% increase in the probability that a locality enters the program. Given the *fuzzy* nature of the discontinuity, our identification yields a local intent to treat effect.⁴ No evidence of sorting of localities around the discontinuity was found based on pre-existing observables.

Our first set of results shows that program eligibility indeed expanded healthcare utilization at public facilities, with an average increase of around 12% in outpatient care visits. We also show that this increase in formal, professional healthcare is accompanied by a decline in seeking alternative health services, such as non-Western medicine. This change is driven by both children and women aged 20 to 49, which is consistent with the program requirements.

We then focus on observable healthcare and behaviors related to reproductive health and chronic conditions, showing positive impacts on both access to and utilization of preventive care, such as contraceptives, prenatal doctor visits, and screenings for chronic diseases and cancer. While some of these results are mechanical due to the program requirements (see online appendix Table A1), others —such as taking contraceptives and undergoing screenings for chronic conditions— were not explicitly targeted by the program.

Our last set of outcomes explore whether this increased utilization led to congestion and, consequently, to crowding-out of certain profiles of users at these public clinics. We find evidence of an increase in waiting times and a decline in self-reported patient satisfaction. These findings are consistent with a decline in some dimensions of quality of care. Nonetheless, the demographic

⁴*Complier* localities are those that receive the PROGRESA cash transfer when they are eligible (i.e. to the right of the discontinuity) and would not receive it if they are not eligible (i.e. to the left of the regional cutoff). As such, we cannot speak to the effects for always-takers (i.e., localities that regardless of their poverty index would always find a way to get the program) or never-takers.

composition of users does not suggest a reduction in attendance of elderly people, who are not targeted through this program. Moreover, the data does not allow us to identify any potential adverse health impacts stemming from congestion.

Taken together, our results suggest that embedding preventive healthcare in a cash transfer can generate positive impacts in utilization, not just on targeted items but across other dimensions as well. However, inducing higher utilization without the adequate supply expansions may hinder some of these efforts. Note as well that we cannot speak to the importance of conditioning the cash transfer since we do not observe variation along this dimension in our context. Nevertheless, given the ubiquity of CCTs in other contexts, our findings suggest a particular way in which uptake of preventive healthcare may be increased.

Our paper contributes to different strands of the literature. First, given the considerable surge in demand at the local clinics, our paper adds to work that has analyzed how changes in administrative rules that impose or suggest health requirements impact health outcomes. Studies for the US have shown that vaccine mandates for school enrollment were essential drivers for eradicating transmissible diseases among children (Lawler, 2017) and that the recent push for exemptions has led to adverse health effects (Hair et al., 2021). Other studies have focused on guidelines issued by governments and non-profit organizations. Einav et al. (2020) employ a change in the age screening requirement for mammograms and show that using RCTs to measure compliers' effects on mammogram recommendation might significantly underestimate the benefits of screening.

Second, we advance the literature on the health impacts that result from CCT program designs. The latter has been studied from different angles. Some papers have analyzed the impact of implementing a CCT versus not.⁵ They have confirmed increased demand resulting from these designs and analyzed specific outcomes influenced by the program. Analyzing a policy in Tanzania, Evans et al. (2019) find an initial surge in demand which dissipated after 2.5 years, and an increase in the likelihood of seeking treatment when ill. For a CCT in Nicaragua, Barham and Maluccio (2009) show that the program managed to increase vaccination rates and successfully reached levels required to eradicate certain diseases, an achievement that could not be met using other interventions. For the case of PROGRESA, Barham and Rowberry (2013) use the program's phase-

⁵Other work has focused on the contribution of the health conditionality itself. Akresh et al. (2013) do so by comparing conditional and unconditional cash transfer programs. Attanasio et al. (2015) compare children born before and after their mother registered to a Colombian CCT program.

in to find a reduction in elderly mortality, mainly driven by impacts on diabetes and infectious diseases. In the same context, [Barham \(2011\)](#) also finds a decline in rural infant mortality associated with the program. A review of additional literature is provided by [Lagarde et al. \(2007\)](#), generally supporting that CCTs are successful at promoting preventive health and in some cases health outcomes.

Finally, our paper gives evidence of negative externalities and thus contributes to work in the CCT and health policies literature looking at spillovers. [Andrew and Vera-Hernández \(2022\)](#) develop a general equilibrium model and provide evidence of how the sudden increase in demand promoted by a policy that incentivizes demand can harm part of the population. [Guerrero et al. \(2020\)](#) analyze how the Peruvian CCT motivated the substitution from informal to formal health services both for targeted and non-targeted recipients. [Suarez and Maitra \(2021\)](#) looks at positive spillovers in women’s and the elderly’s health outcomes in the Colombian CCT, *Familias en Acción*, likely coming from increased information at the locality level. As for PROGRESA, [Avitabile \(2021\)](#) finds an increase in pap smear screening among ineligible women, but no externalities in non-gender-specific tests, perhaps driven by a weakening of husbands’ opposition to pap smear screening. [Gertler \(2000\)](#) finds that PROGRESA significantly increased utilization of public health clinics for preventive care, including prenatal care, child nutrition monitoring, and adult check-ups. The program also lowered the number of inpatient hospitalizations, which is consistent with the hypothesis that PROGRESA lowered the incidence of severe illness. Moreover, there was no reduction in the utilization of private providers, suggesting that the increase in utilization at public clinics was not substituting public for private care. Lastly, [Hernández et al. \(1999\)](#) find an increase in visits to PROGRESA clinics compared to the rest. This increase ranges from 11.5% to 19.5%, and is greater for age groups 25-44 and 5-14. The authors associate this with the co-responsibility scheme that involves all household members. We extend this literature by focusing on various outcomes of preventive healthcare and by quantifying any potential downsides (in terms of waiting times and self-reported quality of care) from this surge in demand that was unaccompanied by an increase in supply.

The remainder of the paper is organized as follows. The next section provides details on the program. Section 3 describes the data. Section 4 lays out the empirical strategy. Section 5 presents the results. Lastly, Section 6 concludes.

2 Background on PROGRESA

PROGRESA is one of the most well-known CCT programs, backed by its solid institutional foundation and rigorous evaluation design.⁶ Introduced in 1997, the main objective of PROGRESA was to improve the health and development of children, including education and nutrition ([Secretaría de Desarrollo Social, 2000](#)). A secondary objective was to improve adult health ([Barham and Rowberry, 2013](#)). PROGRESA sought to promote health care for all family members via a predominantly preventive approach ([Secretaría de Desarrollo Social, 2000](#)). Before its expansion in 2004, PROGRESA targeted poor rural localities only.⁷ By 2001, the program had already been extended to 67,539 localities, which amounts to a third of all localities in Mexico. By 2004, the program changed its expansion design, opening to urban areas and reaching a total of 82,973 localities.

Before 2004, eligibility for PROGRESA was described in the program’s documentation as a multi-stage process. First, localities lacking access to schooling and healthcare infrastructure were not made eligible due to the impossibility of verifying the school and health attendance requirements established by the program ([Hernández Licona et al., 2019](#)). Second, the remaining localities were ranked based on a poverty index constructed from indicators collected during the 1990 census and the 1995 short census. Government officials established regional cutoff values for program eligibility, which is the source of variation that we exploit and further describe below. Third, once a locality was eligible, detailed information (including demographic characteristics and durable asset ownership) was gathered from every household in the locality to determine whether each particular household would become a program recipient.⁸ Precise eligibility factors were not known by beneficiaries nor local authorities in order to avoid strategic manipulation ([Aguilar and Vicarelli, 2022](#)).

Household cash transfers were conditioned on children’s school enrollment, preventive healthcare visits for all household members, and participation in health education training sessions to be

⁶The design of PROGRESA included a randomized trial implemented between 1997 and 2000, with follow-up surveys in 2003 and 2007 to assess its short and medium-term benefits ([Skoufias, 2001](#); [Behrman et al., 2005](#)). Eligible households in treatment localities began receiving benefits in 1998, and in 2000, eligible households in control localities joined the program ([Skoufias et al., 1999](#); [Coady, 2000](#); [Gertler and Fernald, 2004](#)). More recent studies have also evaluated the long-run effects of the program ([Aguilar et al., 2019](#); [Parker and Vogl, 2021](#)).

⁷Localities are the smallest administrative unit in Mexico, which are, in turn, grouped into municipalities. The 2000 census recorded 199,391 localities in 2,445 municipalities. Rural localities are defined as those below 2,500 inhabitants. However, this restriction was not strictly implemented in practice, although the program was indeed focused on smaller and less developed localities.

⁸Additionally, a fourth step consisted of a verification done by a local council, which would vouch if eligible households were in actual need of the cash transfer. Qualitative evidence suggests that this step rarely influenced the final list of recipients.

attended by at least one member (Barham and Rowberry, 2013). Online appendix Table A1 shows the mandatory number of health check-ups per year by different age groups. In addition, pregnant women were required to have five check-ups and two additional ones while breastfeeding (Secretaría de Desarrollo Social, 2000). Cash transfers were delivered to the female head of the household every two months. Compliance was fostered by providing an appointment book to beneficiaries (Barham, 2011).

Under PROGRESA, public health clinics were required to provide a package of services (see Table A2). This included: family planning, education on basic sanitation and accident prevention, prenatal, delivery, and postpartum care, child growth monitoring, vaccination, anti-parasitic treatments, and prevention and treatment of diarrhoea, respiratory infections, tuberculosis, high blood pressure, and diabetes (Barham and Rowberry, 2013). PROGRESA had the support of 10,141 clinics, in which almost 40,000 institutional service providers collaborated, including 12,787 physicians (1.3 per clinic on average) and more than 14 thousand nurses (1.4 per clinic). Additionally, it had 23,830 health assistants (2.3 per clinic) in charge of health promotion and prevention activities (Hernández et al., 1999). PROGRESA focused exclusively on primary health care services (Secretaría de Desarrollo Social, 2000). Importantly, the expansion of PROGRESA was not accompanied by new healthcare infrastructure (see Figure A7 and Table A9).

3 Data

We combine information on program roll-out with healthcare utilization outcomes from various administrative records and a national health survey. This methodology allows us to construct a locality-level data set for our primary empirical analysis. This section describes each data source in detail.

PROGRESA Administrative Records. We obtained access to data detailing the number of cash transfers paid out at each locality per year. With this information we know how the program roll-out occurred by looking at the first payment at each locality. From PROGRESA we also obtain the geographical organization of the program by regions. We add the poverty index (or marginality index) calculated by the Federal government agency CONAPO (National Population Council) in

1995. Note that a *higher* index value corresponds to a *lower* socioeconomic status. This continuous index is the result of a principal components analysis that uses as inputs variables from the 1990 census and 1995 short census (Skoufias et al., 1999). Altogether, these data allow us to infer the regional poverty index cutoffs for inclusion into the program, which we will employ in a regression discontinuity design.

National Health Survey (ENSA). The 2000 wave of the ENSA is a nationally representative survey of the population’s health status and healthcare utilization. It is an important public policy tool that disseminates the health status and nutritional conditions of various population groups in Mexico (Gutierrez et al., 2012). ENSA collects information on households at the individual level and has age-specific questionnaires for adults aged 20 or older, teenagers aged 10 to 19, and children under ten. In addition, it contains specific questions for all household members that utilized health services during the year prior to the survey. We construct various measures from this publicly available survey.

First, we generate indicators for medical care if at least one person in the household was sick during the two weeks prior to the survey. Medical care information includes: whether the person was attended by a physician in a clinic, whether the doctor prescribed medications, and whether the person self-medicated. To complement this information, an indicator for whether anyone in the household used healthcare services during the last 12 months is available. For young children aged four and under, we observe if they received medical attention conditional on being sick within the last two weeks. Lastly, we construct a self-reported indicator for whether a person’s health improved after receiving medical care. We also break down some of these utilization metrics by age groups.

Second, we focus on health behaviors related to reproductive health and chronic diseases. For the former, we measure whether individuals aged 12 and over received family planning information, whether pregnant women received prenatal information, whether adults used contraceptive methods, the number of prenatal care visits, whether expectant mothers received prenatal care starting in their first trimester, and whether women have ever been pregnant since 1994 and during the year 2000. For chronic conditions, we consider indicators for having a diabetes and/or high blood

pressure (HBP) diagnosis, whether patients are on a prevention program for diabetes and/or HBP, and indicators for whether adult women received a pap smear test or mammogram.

Lastly, we compile information on proxies for self-reported quality of care. We consider waiting times, a dummy indicating if the waiting time was short,⁹ and a dummy indicating if patients consider that the time spent with the doctor was enough.

Public clinics’ administrative data. We complement the survey with utilization data from administrative records of public clinics. These data are collected by the Ministry of Health. The information available is at the clinic level from the years 2000 through 2003. The data allow us to observe consultations, excluding emergency room visits, which we aggregate up to the yearly level. Although all public clinics are required by law to report this information, we consider only facilities belonging to the Ministry of Health since PROGRESA recipients were more likely to utilize services at clinics from this institution.¹⁰ We use geographic identifiers to locate the clinic and associate its information with localities.

National System of Health Quality (INDICAS). This system is maintained by the Ministry of Health and is designed to track different measures of quality of care at public clinics. We use information for 2003 since data for previous years is unavailable. The 2003 records include information from 3,794 sampled clinics. To the best of our knowledge, these clinics are randomly sampled by the Ministry of Health to be representative of the quality of care throughout the public system. From INDICAS, we obtain average waiting time reported by the clinic’s personnel, the share of users satisfied with waiting times, whether any complaints were filed, and the share of patients that received an adequate explanation of their health from the doctor.¹¹

⁹The benchmark for a standard wait time is at most 50 minutes for rural public clinics in Mexico (Ruelas et al., 2002).

¹⁰The Mexican Institute for Social Security (IMSS) provides healthcare for formal workers and their families. The Civil Service Social Security and Services Institute (ISSSTE) provides care for government employees and their families. The target population of PROGRESA is unlikely to be eligible for care at facilities run by either of these public institutions. Results hold but are noisier if we include all public clinics in our analysis.

¹¹In order to collect this information, the Ministry of Health trained personnel to capture the survey information from the sampled clinics. The reporting frequency is every two months. Surveyors report on waiting times obtained from clinic staff and sample a subset of patients for questions about satisfaction with waiting times and whether the doctor explained their health status to them. Surveyors are also asked to log complaints.

Table 1 presents descriptive statistics. Each panel corresponds to a different set of variables. Columns indicate the sample (based on available responses), the average, the standard deviation, and the data set from which we obtain each variable.

Table 1: **Descriptive Statistics**

Variable	Description	N	Mean	SD	Data set
Panel A: First Stage					
1995 marginality index	Locality poverty threshold to receive treatment	934	-0.939	0.936	ENSA
Treatment 2000 (0,1)	Locality dummy if entered program after 1999	934	0.480	0.500	ENSA
1995 marginality index	Locality poverty threshold to receive treatment	10608	-0.608	0.920	Admin. data
Treatment 2000 (0,1)	Locality dummy if entered program after 1999	10608	0.625	0.484	Admin. data
Panel B: Use of health services					
Total visits	Total medical visits in 2000 (national)	14373	837.1	1512	Admin. data
Child received treatment (0,1)	Child dummy if received doctor treatment	10460	0.248	0.432	ENSA (minors)
Health services user (0,1)	HH dummy if received attention last year	174566	0.287	0.453	ENSA (households)
Self-medication (0,1)	User dummy if self-medication	174092	0.0210	0.143	ENSA (households)
Health outcome (0,1)	User D if health improved after consultation	80873	0.648	0.478	ENSA (users)
Panel C: Use of health services by ages					
Users aged 0-4 yo (0,1)	HH dummy if utilized services and aged 0-4 yo	19700	0.395	0.489	ENSA (households)
Users aged 5-14 yo (0,1)	HH dummy if utilized services and aged 5-14 yo	41646	0.243	0.429	ENSA (households)
Users aged 15-19 yo (0,1)	HH dummy if utilized services and aged 15-19 yo	16970	0.196	0.397	ENSA (households)
Users aged 20-49 yo (0,1)	HH dummy if utilized services and aged 20-49 yo	68965	0.262	0.440	ENSA (households)
Users aged 50-64 yo (0,1)	HH dummy if utilized services and aged 50-64 yo	16935	0.373	0.484	ENSA (households)
Users aged 65+ yo (0,1)	HH dummy if utilized services and aged 65+ yo	10208	0.445	0.497	ENSA (households)
Panel D: Reproductive health					
Contraceptives (0,1)	Dummy if adult uses effective contraceptives	22914	0.481	0.500	ENSA (adults)
Prenatal check-ups	Number of pregnancy check-ups	10846	6.249	3.771	ENSA (adults & teens)
Has been pregnant (0,1)	D if woman/teen has been pregnant since 1994	20781	0.527	0.499	ENSA (adults & teens)
Recent pregnancy (0,1)	Dummy if woman/teen was pregnant in 2000	24104	0.0664	0.249	ENSA (adults & teens)
Panel E: Chronic disease and prevention					
Diabetes (0,1)	D if adult has been diagnosed with diabetes	40517	0.0666	0.249	ENSA (adults)
High blood pressure (0,1)	D if adult diagnosed with HBP	34647	0.173	0.378	ENSA (adults)
Pap smear test (0,1)	Dummy for preventive pap smear testing	52537	0.105	0.307	ENSA (households)
Breast cancer (0,1)	Dummy for preventive breast cancer testing	27431	0.106	0.308	ENSA (adults)
Panel F: Service characteristics					
Waited 50- minutes (0,1)	D if user-reported waiting was below 50 mins	79350	0.718	0.450	ENSA (users)
Wait satisfaction	Share of users satisfied with waiting time	2375	87.30	16.81	INDICAS
Complaint (0,1)	Dummy if there was a user complaint	2375	0.086	0.281	INDICAS
Diagnosis explained	% of users to which doctor explained health status	2375	93.64	13.99	INDICAS

Note: (0,1) denotes a dummy variable. Statistics are reported for 2000, except INDICAS which reports information for 2003.

HH: Household. D: Dummy. yo: years old.

4 Empirical Strategy

We focus our analysis on the pre-2004 rural expansion of the program. Following the expansion strategy and rules described in Section 2, we employ a regression discontinuity design, since localities were added to the program based on their poverty index and specific regional cutoffs.

4.1 Eligibility for PROGRESA

For purposes of this program, the country was partitioned into 41 regions.¹² As outlined before, the federal government established different thresholds of the locality poverty index for program eligibility that varied by region and time. Whenever PROGRESA was expanded (which occurred almost every year before 2004), localities with an index value above the cutoff were determined as eligible. Although we were unable to retrieve official documents containing precise details of the regional cutoffs, we know from official documentation that the key input to determine eligibility was the 1995 poverty index provided by CONAPO ([Secretaría de Desarrollo Social, 2000](#)).

Using the fact that administrative records allow us to identify the exact year of enrollment to PROGRESA at the locality level, we follow a data-driven approach to identify the regional thresholds for inclusion into the program. First, we use actual payments to determine the entry year of each locality. That way, at the locality level, we know for each year if the locality is already enrolled in the program. Then, using enrollment as a dependent variable and CONAPO’s 1995 poverty index as the running variable, we follow the standard approach for an RDD estimation by implementing a local linear regression discontinuity (see online appendix [Appendix 2](#) for details). We implement this recursively for different values of the cutoff per year-region. We keep the cutoff value with the maximum discontinuous jump in localities’ enrollment. Overall, this allows us to identify eligibility thresholds for each region.¹³ This approach is broadly adapted from the method to identify tipping points in ([Card et al., 2008](#)). However, we also show robustness to simply implementing their method directly in Table [A14](#).


¹²See online appendix Table [A3](#) for details. These regions do not correspond to the 32 Mexican administrative states. Some regions are located within a single state, but others expand beyond state boundaries. Later iterations of the program condensed these 41 regions into 19.

¹³Previous work analyzing program effects in rural areas has used a single cutoff value of -1.2 ([Alix-Garcia et al., 2013](#)). We compare our first stage results to those produced by using this constant cutoff value in Figure [A2](#). It should be noted that our sample of localities is more extensive than the one considered in [Alix-Garcia et al. \(2013\)](#), which does not allow us to make a direct comparison between their estimates and ours. We replicated their strategy with our sample, which led to an estimated first stage change of 15.82 p.p., close to half of our estimate.

Figure 1 shows the result of this discontinuity identification by pooling together all the regions and re-centering all the cutoffs at zero. We present a binned scatterplot of the share of localities that began receiving PROGRESA in the year 2000 for different bins of the distance to the region-specific threshold. As can be seen, we obtain a strong first stage: the discontinuous change at zero in the percent of localities enrolled is considerable, with an estimated increase of 35%. We contrast this value with the first stage effects from alternative methods for identifying the region-specific cutoffs for program eligibility in Figure A2.

Additionally, we overlay in gray the density of localities by values of the re-centered running variable, showing that it is continuous around the threshold. This suggests that there was no evident strategic modification of the poverty index in order to become eligible for the program. We show a formal test for this in Figure A3.

Figure 1: **First Stage**



Note: This figure shows a scatter plot of the share of localities that have entered the program (left axis) by values of the re-centered poverty index at the locality level, and the density of this re-centered running variable (right axis). The vertical line denotes the minimum value for program eligibility. See Appendix 2 for more details.

4.2 Regression Discontinuity Design

Given the design of the program, a simple comparison between localities that were enrolled in PROGRESA and those that were not could be confounded by unobserved factors correlated with eligibility. To address this, we exploit the discontinuity described before to compare localities marginally included in the program with those just excluded. Although many socioeconomic factors determine eligibility via the poverty index, these variables do not change discontinuously around the threshold.¹⁴ In contrast, the program roll-out does change abruptly, as shown in Figure 1, allowing us to estimate a causal effect of PROGRESA.

Figure 2 shows a map of the proportion of treated localities by municipality using the localities in the ENSA and administrative records samples. There is considerable heterogeneity across space and no clear spatial or regional correlations.

Assuming continuity of other socioeconomic characteristics around the threshold, and given the *fuzzy* nature of the discontinuity, the RDD estimator gives us the local intent to treat (ITT) effect of the PROGRESA CCT on health outcomes. Our parameter of interest (τ_x), corresponds to the local ITT, and is defined as follows:

$$\tau_c = E[(Y_i(1) - Y_i(0))|X_i = c, complier] Pr(complier|X_i = c) \quad (1)$$

In our context, to define a complier, eligibility is determined by the administrative rule described in the previous subsection and treatment as receiving PROGRESA. To estimate τ_x , we follow the state-of-the-art procedure. We will employ the difference of approaching the conditional mean of the outcome from the right and left of the discontinuity, which gives variation resulting from the proportion of treated individuals. We use the following specifications:¹⁵

¹⁴We show evidence supporting the continuity assumption in appendix Figure A4.

¹⁵Following Calonico et al. (2014b), we use the following approach to estimate the ITT. Given the discontinuous change in the proportion of treated localities at the threshold, we will estimate the ITT using the discontinuous change in the expected value of Y at the threshold:

$$\begin{aligned} \tau_x &= \mu_+ - \mu_- \\ \mu_+ &= \lim_{x \rightarrow c^+} E[Y_i|X_i = x] \\ \mu_- &= \lim_{x \rightarrow c^-} E[Y_i|X_i = x] \end{aligned}$$

$$\begin{aligned}
Y_{ijr} &= \alpha_L + f(index_{jr} - c_r) + \varepsilon_{ijr} \\
Y_{ijr} &= \alpha_R + f(index_{jr} - c_r) + \varepsilon_{ijr} \\
\tau_c &= \alpha_R - \alpha_L
\end{aligned} \tag{2}$$

where Y_{ijr} is an outcome for surveyed individual i in locality j and region r (or, alternatively for clinic i in locality j and region r), $index_{jr}$ is CONAPO’s locality-level poverty index, c_r is the region r specific cutoff identified above, $f(\cdot)$ is a flexible function of the running variable, and ε_{ijr} is the idiosyncratic error term. We follow [Calonico et al. \(2014b\)](#) for calculating local polynomial RDD estimators.¹⁶ The first specification employs only observations to the left of the discontinuity (up to the bandwidth) and the second those to the right. We estimate both equations using a triangular kernel, as well as using a local-polynomial of degree one. We specify a fixed bandwidth of 0.5 for all specifications (see [Appendix 4](#) for bandwidth sensitivity analyses).¹⁷

It is important to mention that the ITT estimate, captures the overall effect of the program. We might think that the main driver of the effects found is the conditionality established by the program –of regularly attending the clinic for preventive checkups–, however, we cannot rule out that other components of the program might be partially driving the effects. For instance, the cash transfer component might induce a greater demand for preventive checkups if they are considered a normal good in this context.


To help validate the RDD approach, we show first-stage evidence that the probability of receiving PROGRESA cash transfers increased discontinuously around the eligibility cutoff as discussed above in [Figure 1](#). We also note that we do not find evidence that localities manipulated their score in order to become eligible for the program. A formal test of this ([McCrary, 2008](#)) is shown in [Figure A3](#).

¹⁶We rely on the Stata command developed in [Calonico et al. \(2017\)](#).


¹⁷This bandwidth is similar to the one obtained using the mean-squared error-optimal bandwidths for several of our outcomes across different data sets ([Calonico et al., 2017](#); [Imbens and Kalyanaraman, 2012](#)).

Figure 2: **Treatment per Municipality**

(a) Health services household survey (ENSA 2000)



(b) Administrative data



Note: This figure shows the share of localities that are treated by municipality. The top map considers the sample included in the ENSA survey and the bottom map the sample from administrative records.

5 Results

We present our results in this section by groups of outcomes, drawing on different variables from three data sets. All specifications follow the empirical strategy outlined above and use a bandwidth of 0.5 across all outcomes for consistency.¹⁸ We show robustness to the optimal bandwidth in online appendix Table A13 and present sensitivity analyses on the bandwidth in Appendix 4. For clarity, we present RDD plots showing binned means of the outcome variable around the program eligibility cutoff and a local polynomial of degree one. Shaded areas represent 95% confidence intervals. We also complement this with tables showing the RDD estimates with standard errors robust to heteroskedasticity.

We present a battery of robustness checks. Online appendix Table A10 also considers standard errors clustered at the locality level, with very similar results. Furthermore, we show robustness to multiple hypothesis testing by constructing an index for each set of outcomes in Table A11 and Figure A14 and by using resampling methods to control for the familywise error rate in Table A12. We also show robustness to using an alternative method for identifying the program cutoffs that directly follows Card et al. (2008) in Table A14. While some of these checks are more taxing than others, the general take-aways and patterns hold across specifications.

5.1 Utilization of health services


We begin with utilization measures from administrative records corresponding to public clinics ascribed to the Ministry of Health. We take the log of total yearly visits as our outcome variable and exploit records from multiple years (2000 through 2003). We show plots of each measure against the distance to the threshold for PROGRESA eligibility in panels (a)-(d) of Figure 3. There is a clear increase in utilization precisely at the threshold across samples. Panel A of Table 2 shows the corresponding point estimates, all of which are positive, large, and mostly statistically significant.¹⁹ We cannot reject that effect sizes are the same across years. On average, we find about a 12% increase in total visits, ranging from a 7% increase in 2000 to a 16% increase in 2003.

¹⁸We estimated the optimal bandwidth for each specification and found that 0.5 was a good approximation for standardizing across outcomes.

¹⁹Only the effect for the year 2000 is not significant at conventional levels. However, given the size of the point estimate, we do not consider this to be a precisely estimated zero, but instead a noisily estimated positive impact. Furthermore, we obtain significant effects for this sample under alternative specifications (see Appendix 3).

We then turn our attention to similar measures in the survey data in panels (e)-(h) of Figure 3 and show the corresponding point estimates in panel B of Table 2. Overall, we obtain positive and significant effects showing that the CCT program led to increased health services utilization. We see an increase in children seeing a doctor when sick, households using health services in general, and health improvement. We also see a decline in the probability of self-medicating. Online appendix Figure A6 and Table A8 show complementary results indicating that medical care and prescriptions from formal doctors increased while utilization of alternative health services (such as non-professional medical staff or non-Western medicine) was either unchanged or decreased.

Figure 3: **Use of health services**



Note: These figures show RDD plots for our outcomes of interest. We present binned means of the outcome variables around the program eligibility cutoff and a local polynomial of degree one. Panels (a)-(d) refer to administrative census data. Panels (e)-(j) consider ENSA 2000 sample data.


Table 2: Use of health services

Panel A: Use of health services (Administrative census data)				
	ln(Total visits)			
Year	2000	2001	2002	2003
$\hat{\tau}_c$	0.0747 (0.0612)	0.1293 (0.0613)	0.1142 (0.0631)	0.1578 (0.0625)
Observations	8926	8977	8594	9444
P-value	0.2220	0.0351	0.0706	0.0115
Mean dependent variable	6.7640	6.7290	6.7620	6.7400
Mean dep. var. left of cutoff	6.7070	6.6850	6.6610	6.6700
Panel B: Use of health services (ENSA 2000 sample)				
	Child received treatment	Used health services	Self- medication	Health improved
$\hat{\tau}_c$	0.0818 (0.0385)	0.0397 (0.0099)	-0.0085 (0.0035)	0.0664 (0.0158)
Observations	10460	174566	174092	80873
P-value	0.0338	0.0000	0.0136	0.0000
Mean dependent variable	0.2480	0.2870	0.0210	0.6480
Mean dep. var. left of cutoff	0.2210	0.2640	0.0288	0.6240

Note: This table shows RDD estimates of the impact of Progresa eligibility on different outcomes that result from estimating Equation 2. All outcome variables are defined in Table 1. Panel A uses administrative data, while Panel B uses survey data. We report conventional standard errors (SE) robust to heteroskedasticity in parenthesis. Additionally, we include the overall mean of the dependent variable and its mean for localities just to the left of the cutoff value for eligibility (-0.1, 0).

The PROGRESA transfer was conditioned on different requirements of healthcare visits for different age groups, with a particular target on children, pregnant women, and new mothers. We partition survey respondents into six age brackets to disentangle our effects by age group. We show RDD plots for these age-specific outcomes in Figure 4, and the corresponding estimates in Table 3. We find large and significant increases in healthcare utilization for children (ages 0 to 4 and ages 5 to 14) and adults (ages 20 to 49). The latter is concentrated among women rather than men, which is consistent with the program requirements for pregnant and breastfeeding women (see online appendix Table A6). We find small and insignificant effects for individuals aged 15 to 19 and elderly people aged 65 and over. Lastly, although noisily estimated, we find a large positive point estimate for older adults (ages 50 to 64).

Figure 4: Use of health services by ages



Note: These figures show RDD plots for our outcomes of interest. We present binned means of the outcome variables around the program eligibility cutoff and a local polynomial of degree one.

Table 3: Use of health services by ages

	0-4 years old	5-14 years old	15-19 years old	20-49 years old	50-64 years old	65+ years old
$\hat{\tau}_c$	0.0557 (0.0316)	0.0433 (0.0180)	0.0084 (0.0265)	0.0368 (0.0159)	0.0531 (0.0345)	-0.0081 (0.0437)
Observations	19700	41646	16970	68965	16935	10208
P-value	0.0776	0.0163	0.7520	0.0206	0.1240	0.8530
Mean dependent variable	0.3950	0.2430	0.1960	0.2620	0.3730	0.4450
Mean dep. var. left of cutoff	0.3820	0.2020	0.1720	0.2350	0.3520	0.4960

Note: This table shows RDD estimates of the impact of Progresa eligibility on different outcomes that result from estimating Equation 2. All outcome variables are defined in Table 1. We report conventional standard errors (SE) robust to heteroskedasticity in parenthesis. Additionally, we include the overall mean of the dependent variable and its mean for localities just to the left of the cutoff value for eligibility (-0.1, 0).

5.2 Reproductive health and chronic diseases

Figure 5 reports impacts on a variety of outcomes related to reproductive health, where we mainly find increases in utilization of contraceptives and prenatal doctor visits. We report point estimates in Table 4. There is a large and significant increase in the use of contraceptives (7.5 pp increase on a baseline utilization of 43%). As for services during pregnancy, we find an average increase of 0.55 pregnancy checkups (or equivalently, 10%) in areas just above the eligibility cutoff for PROGRESA. Lastly, we show that fertility rates are not a potential driver of these results as neither the likelihood of being pregnant since 1994 nor the likelihood of being pregnant in the year 2000 changes across the eligibility threshold. Although informational talks and prenatal checkups were official program requirements (see online appendix Table A1), contraceptive use and actual fertility were not explicitly targeted by the program.


Table 4: **Reproductive health**

	Contra- ceptives	Prenatal checkups	Ever pregnant since 1994	Pregnant in 2000
$\hat{\tau}_c$	0.0755 (0.0312)	0.5530 (0.2971)	0.0078 (0.0319)	0.0044 (0.0147)
Observations	22914	10846	20781	24104
P-value	0.0155	0.0627	0.8060	0.7630
MDV	0.4810	6.2490	0.5270	0.0664
MDV left of cutoff	0.4320	5.6510	0.5680	0.0658

Note: This table shows RDD estimates of the impact of Progresa eligibility on different outcomes that result from estimating Equation 2. All outcome variables are defined in Table 1. We report conventional standard errors (SE) robust to heteroskedasticity in parenthesis. Additionally, we include the overall mean of the dependent variable and its mean for localities just to the left of the cutoff value for eligibility (-0.1, 0). MDV = mean dependent variable.

Figure 6 and Table 5 focus on outcomes related to chronic diseases and preventive care. We find that PROGRESA eligibility increases the probability of having been diagnosed with HBP but has no effect on diabetes diagnoses. The latter may relate to the fact that HBP is more prevalent and more widely under-diagnosed than diabetes in Mexico (Campos-Nonato et al., 2018; Basto-Abreu et al., 2020). This increase in diagnoses is therefore consistent with more access to healthcare. We also show positive effects for pap smears and breast cancer examinations. However, these results are noisily estimated, although the plots in Figure 6 are suggestive of positive impacts. None of these outcomes were explicitly included as part of the behaviors or types of care that the program was attempting to change.


Figure 5: **Reproductive health**



Note: These figures show RDD plots for our outcomes of interest. We present binned means of the outcome variables around the program eligibility cutoff and a local polynomial of degree one.

Taken together, the reproductive health and chronic disease estimates suggest that PROGRESA eligibility had a positive impact on access to and utilization of preventive care. Although we cannot observe direct health outcomes related to these variables, prenatal care visits and timely diagnosis of chronic conditions have been linked in the literature to better health outcomes ([Almond and Currie, 2011](#); [Zhao et al., 2013](#); [Oster, 2018](#)).

Figure 6: **Chronic disease and prevention**



Note: These figures show RDD plots for our outcomes of interest. We present binned means of the outcome variables around the program eligibility cutoff and a local polynomial of degree one. HBP: high blood pressure.

Table 5: **Chronic disease and prevention**

	Diabetes diagnostic	HBP diagnostic	Pap smear test	Breast cancer test
$\hat{\tau}_c$	-0.0009 (0.0115)	0.0318 (0.0193)	0.0076 (0.0129)	0.0199 (0.0150)
Observations	40517	34647	52537	27431
P-value	0.9410	0.0989	0.5570	0.1860
Mean dependent variable	0.0666	0.1730	0.1050	0.1060
Mean dep. var. left of cutoff	0.0687	0.1730	0.1070	0.0692

Note: This table shows RDD estimates of the impact of Progresa eligibility on different outcomes that result from estimating Equation 2. All outcome variables are defined in Table 1. We report conventional standard errors (SE) robust to heteroskedasticity in parenthesis. Additionally, we include the overall mean of the dependent variable and its mean for localities just to the left of the cutoff value for eligibility (-0.1, 0). HBP: high blood pressure.

5.3 Quality of care


Our final set of results considers measures or proxies for quality of care and congestion at public health clinics. As shown above, PROGRESA led to an increase in the utilization of health services. However, during this study period, the program did not consider infrastructure expansions, such as building new clinics or increasing medical staffing. Online appendix Figure A7 and Table A9 show no changes in the supply of public clinics around the cutoff, either in terms of the number of public clinics (intensive margin) or in the presence of any public clinic (extensive margin). Therefore, a natural question is whether the sudden increase in utilization driven by the PROGRESA conditionality led to changes in the characteristics of health services provided.

Figure 7 and Table 6 show results using both the health survey data and records from the government’s quality audit system INDICAS. Focusing on the former (first column of Table 6), given the government’s benchmark of a 50-minute wait, we analyze the probability of waiting below this benchmark. We find a significant decline of 3 pp in the probability of waiting less than 50 minutes, from a baseline share of 72%.

Focusing on the government’s audit system, we find evidence of a decline in the share of clinic patients satisfied with the duration of the wait, an increase in registered complaints, and a decline in the share of patients reporting that they received an explanation of their health status from the doctor. Although these effects are not significant, point estimates are quite large: for example, the effect on complaints logged implies a 29% increase from the baseline mean. Moreover, the plots in panels (b)-(d) of Figure 7 suggest that the effects were not zero.

Overall, our estimates are consistent with the PROGRESA incentive having pushed patients toward higher utilization of health services. However, the lack of additional resources for infrastructure then led to a deterioration in the quality of care as measured by waiting times and satisfaction. An important caveat of this result is that we cannot link it to worse health outcomes. Given the potential gains from increasing healthcare utilization (beyond what we can measure here), it does not seem plausible that this additional congestion was, on average, detrimental to patient health. However, our findings suggest that conditioning government programs without expanding resources may lead to congestion, which may have far-reaching impacts depending on the context.

Figure 7: **Service characteristics**



Note: These figures show RDD plots for our outcomes of interest. We present binned means of the outcome variables around the program eligibility cutoff and a local polynomial of degree one.

Table 6: **Service characteristics**

	Waited 0-50 mins.	Satisfaction with waiting time	Complaint was logged	Medical explanation
$\hat{\tau}_c$	-0.0299** (0.0151)	-3.0930 (2.1854)	0.0252 (0.0279)	-1.3520 (1.7860)
Observations	79350	2375	2375	2375
P-value	0.0478	0.1570	0.3670	0.4490
Mean dependent variable	0.7180	87.3000	0.0863	93.6400
Mean dep. var. left of cutoff	0.7240	87.7500	0.0455	95.2200
Data set	ENSA 2000	INDICAS	INDICAS	INDICAS

Note: This table shows RDD estimates of the impact of Progresa eligibility on different outcomes that result from estimating Equation 2. All outcome variables are defined in Table 1. Panel A uses survey data for 2000 and Panel B uses different survey data for 2003. We report conventional standard errors (SE) robust to heteroskedasticity in parenthesis. Additionally, we include the overall mean of the dependent variable and its mean for localities just to the left of the cutoff value for eligibility (-0.1, 0).

6 Conclusion

This paper analyzes how a CCT policy that focused on children’s schooling and health also led to improvements in adult preventive health due to this additional conditionality of the program. Results show that PROGRESA had positive health effects on the targeted population, following a significant increase in the utilization of health services. We estimate important improvements in reproductive health and chronic disease screening for adults. However, we also find evidence that this program increased waiting times and led to declines in proxies for the quality of health services at public clinics. The latter may suggest critical distributional consequences, with welfare losses for the population segment that was regularly using health services prior to the program, such as those with more inelastic demand for healthcare. However, it is unlikely that this welfare loss would offset the large gains associated with the program, not just in healthcare but in education and other dimensions. Nevertheless, our results suggest that conditioning transfers to public services without the appropriate infrastructure expansion—such as building new schools and clinics—may lead to over-congestion and worse outcomes on some dimensions for at least a subset of the population.

To gauge the benefits of such a policy lever, we present a simple back-of-the-envelope calculation focusing on the increase in HBP diagnoses. As in many other contexts, HBP is underdiagnosed in Mexico, with an average of 1.3 undiagnosed cases per diagnosed case ([Villarreal-Ríos et al., 2002](#)). Furthermore, managing HBP cases costs the public health system an estimated 354 million USD per year ([Arredondo and Zuniga, 2006](#)). These costs include the present value of current and future diagnoses, consultations, drugs, hospitalizations, and treatment of complications (nephropathy, nonfatal myocardial infarction, and nonfatal stroke). The costs of HBP management may be offset by savings from avoiding complications from undiagnosed patients.

Our estimates imply that the program induced 1.7 million new HBP diagnoses among adults due to the program’s conditionality. This suggests an increase of USD 17.6 million in costs associated with chronic disease management (checkups and medication) and a decrease of USD 36.4 million in costs associated with complications and hospitalizations of undiagnosed cases. This calculation assumes that an HBP diagnosis reduces the chances of complications by 98% ([Rosas-Peralta et al., 2016](#)). Therefore, even without considering the private value of health or quality of life, these additional diagnoses potentially saved the public health system around USD 18.8 million annually.

Based on our estimated increase in HBP diagnoses, a prevention effectiveness of 60% would balance out the increased cost of disease management with the savings induced by decreased hospitalizations and treatment of complications, implying that these new diagnoses almost surely were cost-effective.

Since our identification relies on reduced-form local average effects around the cutoffs and given the data availability, we cannot quantify the distribution of welfare consequences in this population. Future work may try to shed light on this issue and identify mediating factors that may have improved the effects on preventive health behaviors or tempered the negative impacts of congestion. Furthermore, we rely on rich but broad administrative and survey data that does not allow us to observe more nuanced outcomes from health visits, such as doctors' treatment choices, nor can we follow the same patients over time. A more detailed dataset may allow for additional insights into how providers adjusted to this inflow of patients and how that may have further impacted health outcomes.

References

- Abrevaya, J. and K. Mulligan (2011). Effectiveness of state-level vaccination mandates: evidence from the varicella vaccine. *Journal of health economics* 30(5), 966–976.
- Aguilar, A. and M. Vicarelli (2022). El Niño and children: Medium-term effects of early-life weather shocks on cognitive and health outcomes. *World Development* 150, 105690.
- Aguilar, A. A., C. Barnard, and G. De Giorgi (2019). Long-Term Effects of PROSPERA on Welfare. *World Bank Policy Research Working Paper* (9002).
- Akresh, R., D. De Walque, and H. Kazianga (2013). Cash transfers and child schooling: evidence from a randomized evaluation of the role of conditionality. *World Bank Policy Research Working Paper* (6340).
- Alix-Garcia, J., C. McIntosh, K. Sims, and J. Welch (2010, 01). Development and deforestation in mexico: Impacts using the discontinuity in eligibility for oportunidades.
- Alix-Garcia, J., C. McIntosh, K. R. Sims, and J. R. Welch (2013). The ecological footprint of poverty alleviation: evidence from Mexico’s Oportunidades program. *Review of Economics and Statistics* 95(2), 417–435.
- Almond, D. and J. Currie (2011). Killing me softly: The fetal origins hypothesis. *Journal of economic perspectives* 25(3), 153–72.
- Andrew, A. and M. Vera-Hernández (2022, 06). Incentivizing Demand for Supply-Constrained Care: Institutional Birth in India. *The Review of Economics and Statistics*, 1–45.
- Arredondo, A. and A. Zuniga (2006). Epidemiologic changes and economic burden of hypertension in latin america: evidence from mexico. *American journal of Hypertension* 19(6), 553–559.
- Attanasio, O. P., V. Oppedisano, and M. Vera-Hernández (2015). Should cash transfers be conditional? Conditionality, preventive care, and health outcomes. *American Economic Journal: Applied Economics* 7(2), 35–52.
- Avitabile, C. (2021). Spillovers and Social Interaction Effects in the Demand for Preventive Health-care: Evidence from the PROGRESA program. *Journal of Health Economics* 79, 102483.

- Barham, T. (2011). A healthier start: The effect of conditional cash transfers on neonatal and infant mortality in rural Mexico. *Journal of Development Economics* 94(1), 74–85.
- Barham, T. and J. A. Maluccio (2009). Eradicating diseases: The effect of conditional cash transfers on vaccination coverage in rural Nicaragua. *Journal of health economics* 28(3), 611–621.
- Barham, T. and J. Rowberry (2013). Living longer: The effect of the Mexican conditional cash transfer program on elderly mortality. *Journal of Development Economics* 105, 226–236.
- Basto-Abreu, A., T. Barrientos-Gutiérrez, R. Rojas-Martínez, C. A. Aguilar-Salinas, N. López-Olmedo, V. De la Cruz-Góngora, J. Rivera-Dommarco, T. Shamah-Levy, M. Romero-Martínez, S. Barquera, et al. (2020). Prevalencia de diabetes y descontrol glucémico en México: resultados de la Ensanut 2016. *Salud Pública de México* 62(1), 50–59.
- Behrman, J. R., P. Sengupta, and P. Todd (2005). Progressing through PROGRESA: An impact assessment of a school subsidy experiment in rural Mexico. *Economic development and cultural change* 54(1), 237–275.
- Calonico, S., M. D. Cattaneo, M. H. Farrell, and R. Titiunik (2017). rdrobust: Software for regression-discontinuity designs. *The Stata Journal* 17(2), 372–404.
- Calonico, S., M. D. Cattaneo, and R. Titiunik (2014a). Robust data-driven inference in the regression-discontinuity design. *The Stata Journal* 14(4), 909–946.
- Calonico, S., M. D. Cattaneo, and R. Titiunik (2014b). Robust nonparametric confidence intervals for regression-discontinuity designs. *Econometrica* 82(6), 2295–2326.
- Campos-Nonato, I., L. Hernández-Barrera, A. Pedroza-Tobías, C. Medina, and S. Barquera (2018). Hipertensión arterial en adultos mexicanos: prevalencia, diagnóstico y tipo de tratamiento. Ensanut MC 2016. *Salud Pública de México* 60, 233–243.
- Card, D., A. Mas, and J. Rothstein (2008). Tipping and the dynamics of segregation. *The Quarterly Journal of Economics* 123(1), 177–218.
- Cawley, J. and J. A. Price (2013). A case study of a workplace wellness program that offers financial incentives for weight loss. *Journal of health economics* 32(5), 794–803.

- Clarke, D., J. P. Romano, and M. Wolf (2020). The romano–wolf multiple-hypothesis correction in stata. *The Stata Journal* 20(4), 812–843.
- Coady, D. P. (2000). The application of social cost-benefit analysis to the evaluation of Progres; final report. Technical report.
- Dupas, P. and E. Miguel (2017). Impacts and determinants of health levels in low-income countries. In *Handbook of economic field experiments*, Volume 2, pp. 3–93. Elsevier.
- Einav, L., A. Finkelstein, T. Oostrom, A. Ostriker, and H. Williams (2020). Screening and selection: The case of mammograms. *American Economic Review* 110(12), 3836–70.
- Evans, D. K., B. Holtemeyer, and K. Kosec (2019). Cash transfers and health: evidence from Tanzania. *The World Bank Economic Review* 33(2), 394–412.
- Gertler, P. (2000). Final report: The impact of progres; on health. *International Food Policy Research Institute, Washington, DC* 35.
- Gertler, P. and L. C. Fernald (2004). The medium term impact of Oportunidades on child development in rural areas. *Unpublished manuscript, University of California at Berkeley*.
- Guerrero, N., O. Molina, and D. Winkelried (2020). Conditional cash transfers, spillovers, and informal health care: Evidence from Peru. *Health Economics* 29(2), 111–122.
- Gutierrez, J. P., J. Rivera-Dommarco, T. Shamah-Levy, S. Villalpando-Hernández, A. Franco, L. Cuevas-Nasu, M. Romero-Martínez, and M. Hernández-Ávila (2012). Encuesta nacional de salud y nutrición 2012. *Resultados Nacionales. Cuernavaca, México: Instituto Nacional de Salud Pública (MX) 2012*, 1–200.
- Hair, N. L., A. Gruber, and C. Urban (2021). Personal belief exemptions for school-entry vaccinations, vaccination rates, and academic achievement. *Journal of Health Economics* 78, 102464.
- Hernández Licona, G., T. De la Garza, J. Zamudio, and I. Yaschine (2019). El Progres; Oportunidades-Prospera, a 20 años de su creación. *Ciudad de México: CONEVAL*.
- Hernández, D., J. G. de León, and G. Vásquez (1999). Más oportunidades para las familias pobres: evaluación de resultados del programa de educación, salud y alimentación: primeros avances.

- Imbens, G. and K. Kalyanaraman (2012). Optimal bandwidth choice for the regression discontinuity estimator. *The Review of economic studies* 79(3), 933–959.
- Jones, D., D. Molitor, and J. Reif (2019). What do workplace wellness programs do? evidence from the illinois workplace wellness study. *The Quarterly Journal of Economics* 134(4), 1747–1791.
- Lagarde, M., A. Haines, and N. Palmer (2007). Conditional cash transfers for improving uptake of health interventions in low- and middle-income countries: a systematic review. *JAMA* 298(16), 1900–10.
- Lawler, E. C. (2017). Effectiveness of vaccination recommendations versus mandates: Evidence from the hepatitis a vaccine. *Journal of health economics* 52, 45–62.
- Levy, D. and J. Ohls (2010). Evaluation of jamaica’s path conditional cash transfer programme. *Journal of Development Effectiveness* 2(4), 421–441.
- McCrary, J. (2008). Manipulation of the running variable in the regression discontinuity design: A density test. *Journal of econometrics* 142(2), 698–714.
- Oster, E. (2018). Diabetes and diet: Purchasing behavior change in response to health information. *American Economic Journal: Applied Economics* 10(4), 308–48.
- Parker, S. W. and T. Vogl (2021). Do conditional cash transfers improve economic outcomes in the next generation? Evidence from Mexico. Technical report, National Bureau of Economic Research.
- Rosas-Peralta, M., G. Borrayo-Sánchez, A. Madrid-Miller, E. Ramírez-Arias, and G. Pérez-Rodríguez (2016). Cardiovascular complications of hypertensive crisis. *Revista Medica Del Instituto Mexicano Del Seguro Social* 54(S1), 67–74.
- Ruelas, B., H. Muñoz, O. Rojas, I. Guillermo, T. Romero, et al. (2002). Lineamientos operativos del sistema de monitoreo para la gestión de la calidad. Primer Nivel. *Secretaría de Salud*.
- Secretaría de Desarrollo Social (2000). Reglas de Operación del Programa de Educación, Salud y Alimentación (PROGRESA).

- Skoufias, E. (2001). Progresa and its impact on the human capital and welfare of household in rural Mexico: A synthesis of the results of an evaluation by IFPRI. Technical report.
- Skoufias, E., B. Davis, and J. R. Behrman (1999). An evaluation of the selection of beneficiary households in the education, health, and nutrition program (PROGRESA) of Mexico. *International Food Policy Research Institute, Washington, DC*.
- Suarez, D. C. and P. Maitra (2021, July). Health spillover effects of a conditional cash transfer program. *Journal of Population Economics* 34(3), 893–928.
- Villarreal-Ríos, E., A. Mathew-Quiroz, M. E. Garza-Elizondo, G. Núñez-Rocha, A. M. Salinas-Martínez, and M. Gallegos-Handal (2002). Costo de la atención de la hipertensión arterial y su impacto en el presupuesto destinado a la salud en México. *Salud pública de México* 44(1), 7–13.
- Zhao, M., Y. Konishi, and P. Glewwe (2013). Does information on health status lead to a healthier lifestyle? Evidence from China on the effect of hypertension diagnosis on food consumption. *Journal of health economics* 32(2), 367–385.

Online Appendices

Appendix 1 Additional Details on the Program

Table A1: Required health check-ups and actions under Progresa

Panel A: Children		
Age group	Check-ups frequency	Actions
less than 4 months	3 check-ups: At 7 and 28 days	Vaccines and growth control weight and height control
4 to 24 months	8 check-ups: At 4, 6, 9, 12, 15, 18 21, and 24 months. Additionally, 1 monthly weight and height check-up.	Vaccines and development control Weight and height control Nutrition monitoring Early disease detection
2 to 4 years	3 yearly check-ups: Once every 4 months.	Vaccines and growth control Weight and height control Deworming Early disease detection
5 to 16 years	2 yearly check-ups: Once every 6 months.	Vaccines and growth monitoring Early disease detection
Panel B: Women		
Group	Check-ups frequency	Actions
Pregnant	5 prenatal check-ups	Nutrition orientation Pregnancy development monitoring Iron and tetanus toxoid provision
Puerperium and lactation	2 check-ups: 1 immediately after childbirth 1 during lactation period	Family planning talks Nutrition orientation Newborn care talks Breast feeding promotion talks
Panel C: Teens and adults		
Age group	Check-ups frequency	Actions
17 to 60 years	1 yearly check-up	Reproductive health talks Family planning talks Early disease detection
More than 60 years	1 yearly check-up	Early detection of chronic disease and neoplasm

Source: [Secretaría de Desarrollo Social \(2000\)](#).

Table A2: **Basic package of health services under Progresa**

Basic sanitation at family level
Family planning
Prenatal, childbirth, puerperium, and newborn care
Surveillance of child nutrition and growth
Immunizations
Management of diarrhea cases at home
Antiparasitic treatment for families
Management of acute respiratory infections
Prevention and control of pulmonary tuberculosis
Prevention and control of arterial hypertension and diabetes mellitus
Prevention of accidents and initial management of injuries
Community training for health self-care
Detection and control of cervical cancer

Source: [Secretaría de Desarrollo Social \(2000\)](#).

Appendix 2 Program Eligibility Threshold

PROGRESA classifies localities into 41 regions (see Table A3). Government agency CONAPO provides the 1995 poverty index by locality. We identify the regional poverty threshold eligibility for the program directly from the data, following a method inspired by Card et al. (2008). Specifically, for a given region r , we estimate the following equation:

$$Enrolled_{jr} = \beta_0^s + \beta_1^s(index_{jr} - c_r) + \epsilon_{jr} \quad (A1)$$

where $Enrolled_{jr}$ is a dummy indicating if locality j in region r is enrolled in the program, $index_{jr}$ is the CONAPO 1995 poverty index for locality j in region r , c_r denotes a fixed value of the poverty index, and ϵ_{jr} is an idiosyncratic error term.

In this specification, $s = \{l, r\}$, where l and r stand for left and right with respect to the discontinuity c_r . Essentially, we estimate (A1) twice: one with localities with $index_{jr} \in [c_r - h, c_r)$ and a second for localities with $index_{jr} \in [c_r, c_r + h]$. A triangular kernel is used to weight observations, giving a larger importance to localities with $index = c_r$. With this two estimations, the discontinuous jump of proportion of localities enrolled in the program is estimated as $\theta = \beta_0^r - \beta_0^l$.

We recursively estimate θ by changing the value of c_r by region-year. We then compare the estimates of θ and identify the cutoff as the value c_r for which θ is largest, conditional on being significant at the 5% significance level.

Figure A1 shows two examples of our estimated cutoffs via this algorithm. The first plot denotes a fuzzy threshold and the second plot shows a very clean and sharp threshold.


We consider two alternatives for identifying the region-specific thresholds for program eligibility. First, we follow Alix-Garcia et al. (2013) and simply use the value of -1.2 for all regions. Second, we closely follow the method for identifying tipping points as developed in Card et al. (2008). This involves estimating a quartic polynomial on either side of a candidate threshold k . The cutoff is identified as the one that yields the largest positive jump from the left to right-side of the threshold. This method requires that the researchers focus their search area on a given range of potential threshold values $[k_0, k_1]$, in order to avoid outliers. We use the -1.2 threshold in Alix-Garcia et al. (2010) as a reference point for this range.

Table A4 compares the threshold values under each of the three methods. Figure A2 shows the first stage under each method for the year 2000. For our preferred method of estimated thresholds described above, we estimate a first stage effect of 35%. Using the constant threshold value of -1.2, we obtain an effect of 16%. Lastly, following the tipping points method, we estimate a 29% increase in the probability of receiving the program. Hence, our method for identifying the cutoff values is a significant improvement over the constant threshold criterion.

Focusing on our method for estimating the cutoffs, we further show in Figure A3 that the density of localities is smooth across the cutoff. We calculate a McCrary test statistic of -1.37 with a p-value of 0.17.

Finally, Figure A4 shows continuity tests using variables from the 1990 Census. We show that labor force participation, share of population at school, share of households with piped water, and share of households without a dirt floor are all continuous around the program eligibility threshold, further lending credibility to our identification strategy.

Figure A1: **Examples of Regional Cutoffs**




Note: Authors' calculations with data from Progresa and Conapo.

Table A3: Progreso Regions

Region	States
1	Chiapas
2	Chiapas
3	Oaxaca, Puebla, Veracruz
4	Hidalgo, Puebla, Tlaxcala
5	Guanajuato, Hidalgo, Querétaro, San Luis Potosí
6	Guerrero
7	Chihuahua, Durango
8	Coahuila, Durango, Nuevo León, Zacatecas
9	Campeche
10	Guerrero, Oaxaca
11	Campeche, Quintana Roo
12	San Luis Potosí
13	Oaxaca
14	Guanajuato
15	Chiapas
16	Oaxaca
27	Guanajuato, Guerrero, Edomex, Michoacán
28	San Luis Potosí
31	Sinaloa, Sonora
32	Tamaulipas
33	Jalisco, Nayarit
34	Jalisco
35	Jalisco, Michoacán
36	Colima, Jalisco , Michoacán
37	Yucatán
38	Morelos
39	Tabasco
40	Tabasco
41	Oaxaca, Veracruz
42	Veracruz
43	Jalisco
44	Puebla
45	Baja California
46	Baja California Sur
47	Coahuila
48	Tamaulipas
49	Sonora
50	Oaxaca
51	Veracruz
52	Aguascalientes
53	Edomex

Note: Regions 17 to 26, 29 and 30 denote an old classification no longer in place during the study period.

Figure A2: First Stage sensibility to the poverty threshold identification strategy




Note: These figures show a scatter plot of the share of localities that have entered the program (left axis) by values of the re-centered poverty index at the locality level. The vertical line denotes the minimum value for program eligibility. We do not restrict the sample. The first plot shows the first stage using our preferred method as in the main text, the second plot uses the constant threshold value of -1.2 as in [Alix-Garcia et al. \(2010\)](#), and the last plot follows the method in [Card et al. \(2008\)](#) by calculating quartic polynomials.

Table A4: **Region-specific eligibility thresholds by method**


Region	Estimated Cutoff	Constant cutoff	Tipping point cutoff
1	-1.2	-1.2	-1.19
2	-0.6	-1.2	-0.89
3	-0.95	-1.2	-0.99
4	-1.2	-1.2	-1.19
5	-0.95	-1.2	-0.99
6	-1.2	-1.2	-1.09
7	-1.2	-1.2	-1.19
8	-1.2	-1.2	-1.19
9	-1.4	-1.2	-1.01
10	-1.05	-1.2	-1.1
11	-1.2	-1.2	-1.17
13	-0.95	-1.2	-0.89
14	-0.95	-1.2	-0.99
15	-1.15	-1.2	-1.18
16	-1.4	-1.2	-0.95
27	-1.05	-1.2	-0.99
28	-1.2	-1.2	-1.22
31	-0.95	-1.2	-0.99
32	-1.2	-1.2	-1.09
33	-1.2	-1.2	-1.21
34	-0.95	-1.2	-1.01
35	-0.95	-1.2	-0.99
36	-1.1	-1.2	-0.99
37	-1.3	-1.2	-1.38
38	-0.95	-1.2	-0.99
40	-1	-1.2	-0.99
41	-0.95	-1.2	-0.99
42	-0.95	-1.2	-1
43	-1.05	-1.2	-0.99
44	-1.15	-1.2	-1.19
45	-0.9	-1.2	-0.9
46	-1.2	-1.2	-1.2
47	-0.95	-1.2	-0.91
48	-1.2	-1.2	-0.87
49	-0.9	-1.2	-1.3
50	-0.95	-1.2	-0.99
51	-0.95	-1.2	-0.9
52	-0.95	-1.2	-0.99
53	-0.8	-1.2	-0.78

This table lists the estimated cutoffs by region for different methods. The first column shows our preferred method as in the main text. The second column shows the constant threshold as in [Alix-Garcia et al. \(2010\)](#). The last column follows the tipping points method in [Card et al. \(2008\)](#) and involves estimating a quartic polynomial on either side of a potential cutoff k , and identifying the largest positive jump within a small window around -1.2 (this number is taken from the cutoff used by [Alix-Garcia et al. \(2010\)](#); this method ignores distances far from this value to address issues with outliers).

Figure A3: **McCrary Manipulation Test**

Note: Authors' calculations with data from Progresa and Conapo.


Figure A4: Continuity test with 1990 data



Note: Authors' calculations with Progresa and INEGI's 1990 Population Census data. These figures show RDD plots for health-associated outcomes in 1990, before Progresa was implemented. We present binned means of the outcome variables around the program eligibility cutoff and a local polynomial of degree one. More indicators were tested using the 1990 census data, showing a continuous behavior around the discontinuity.

Appendix 3 Additional Results

Figure A5: Use of health services: Robustness to dropping clinics with potentially noisy data



Note: Authors' calculations with data from Progresa, Conapo, and the Ministry of Public Health. This figure is analogous to Panels (a)-(d) of Figure 3 in the main text. However, the sample is restricted here to Ministry of Health clinics for which we consistently observe more than 26 weeks of non-zero consultations.

Table A5: **Use of health services: Robustness to dropping clinics with potentially noisy data**

	ln(Total visits)			
	2000	2001	2002	2003
$\hat{\tau}_c$	0.1023 (0.0573)	0.1005 (0.0596)	0.1191 (0.0590)	0.1196 (0.0608)
Observations	8255	8009	7963	7964
P-value	0.0744	0.0917	0.0436	0.0492
Mean dependent variable	6.8570	6.8230	6.8550	6.8430
Mean dep. var. left of cutoff	6.7470	6.7630	6.7120	6.7700

Note: This table is analogous to Panel A of Table 2 in the main text. However, the sample is restricted here to Ministry of Health clinics for which we consistently observe more than 26 weeks of non-zero consultations. This table shows RDD estimates of the impact of Progresa eligibility on different outcomes that result from estimating Equation 2. We report conventional standard errors (SE) robust to heteroskedasticity in parenthesis. Additionally, we include the overall mean of the dependent variable and its mean for localities just to the left of the cutoff value for eligibility (-0.1, 0).

Table A6: **Use of health services by gender (ages 20-49)**

	Use of health services for 20-49 yo		
	All	Female	Male
$\hat{\tau}_c$	0.0368 (0.0159)	0.0517 (0.0224)	0.0181 (0.0216)
Observations	68965	37807	31158
P-value	0.0206	0.0213	0.4030
Mean dependent variable	0.2620	0.3080	0.2060
Mean dep. var. left of cutoff	0.2350	0.2780	0.1820

Note: This table is analogous to Table 3 in the main text. Column (1) replicates the result of Table 3 column (4). Subsequent columns in this table decompose the effect by gender. This table shows RDD estimates of the impact of Progresa eligibility on different outcomes that result from estimating Equation 2. We report conventional standard errors (SE) robust to heteroskedasticity in parenthesis. Additionally, we include the overall mean of the dependent variable and its mean for localities just to the left of the cutoff value for eligibility (-0.1, 0).

Table A7: **Additional Descriptive Statistics on Health Service Utilization**

Variable	Description	N	Mean	SD	Data set
Panel A: Medical attention when sick					
Treated by doctor (0,1)	User dummy if treated by doctor	174346	0.0836	0.277	ENSA (households)
Professional non-doctor (0,1)	Dentist or nurse	174346	0.00147	0.0383	ENSA (households)
Non-Western/alternative doctor (0,1)	Homeopathy, healer, pharmacist	174346	0.00254	0.0503	ENSA (households)
Non-professional (0,1)	Family, friend, community head, other	174346	0.00369	0.0607	ENSA (households)
Panel B: Medication when sick and received treatment					
Medication from doctor (0,1)	User dummy if prescribed by doctor	174092	0.0817	0.274	ENSA (households)
Professional non-doctor (0,1)	Dentist or nurse	174092	0.00142	0.0376	ENSA (households)
Non-Western/alternative doctor (0,1)	Homeopathy, healer, pharmacist	174092	0.00453	0.0672	ENSA (households)
Non-professional (0,1)	Family, friend, community head, other	174092	0.0296	0.170	ENSA (households)
Panel C: Utilization of private clinics for all survey respondents					
Private clinics	Preference for private clinics when sick	168797	0.257	0.437	ENSA (households)

Note: (0,1) Denotes a dummy variable. Statistics are reported for the year 2000.

Table A8: **Additional Results on Health Service Utilization**

Panel A: Medical Attention					
	Treated by doctor	Professional non-doctor	Non-Western alternative doctor	Non- professional	Private preference
$\hat{\tau}_c$	0.0132 (0.0062)	-0.0005 (0.0010)	-0.0003 (0.0017)	-0.0003 (0.0020)	0.0158 (0.0131)
Observations	174346	91545	91545	91545	88966
P-value	0.0320	0.6120	0.8700	0.8580	0.2290
Mean dependent variable	0.0836	0.0017	0.0028	0.0038	0.2570
Mean dep. var. left of cutoff	0.0792	0.0013	0.0018	0.0044	0.2230
Panel B: Medication					
	Medication from doctor	Professional non-doctor	Non-Western alternative doctor	Non- professional	
$\hat{\tau}_c$	0.0122 (0.0061)	0.0001 (0.0012)	0.0017 (0.0017)	-0.0099 (0.0057)	
Observations	174092	91407	91407	91407	
P-value	0.0452	0.9120	0.2960	0.0788	
Mean dependent variable	0.0817	0.0016	0.0048	0.0316	
Mean dep. var. left of cutoff	0.0768	0.0013	0.0013	0.0419	


Note: This table shows RD estimates of the impact of Progresa eligibility on different outcomes that result from estimating Equation 2. All outcome variables are defined in Table A7. Panel A in this table deepens on column (1) of Panel B in Table 2. Panel B in this table deepens on column (4) of Panel B in Table 2. We report conventional standard errors (SE) robust to heteroskedasticity in parenthesis. Additionally, we include the overall mean of the dependent variable and its mean for localities just to the left of the cutoff value for eligibility (-0.1, 0).

Table A9: **Supply of health clinics**

	N clinics in 2000	Any clinic in 2000	N clinics in 2003	Any clinic in 2003
$\hat{\tau}_c$	-0.0142 (0.0384)	-0.0178 (0.0270)	0.0229 (0.0391)	0.0029 (0.0229)
Observations	8555	8555	8244	8244
P-value	0.7120	0.5100	0.5580	0.8990
Mean dependent variable	1.0840	0.8360	1.1780	0.8850
Mean dep. var. left of cutoff	0.9220	0.8650	0.9940	0.9060


Note: This table compares the number of clinics in the treated and untreated localities. Odd columns consider the number of clinics (intensive margin). Even columns show an indicator for a non-zero number of clinics (extensive margin).

Figure A6: **Additional Results on Health Service Utilization**



Note: Authors' calculations with data from Progresa, Conapo, and ENSA 2000. This figure expands on Figure 3 in the main text. All outcome variables are defined in Table A7.

Figure A7: **Supply of health clinics**




Note: Authors' calculations with data from the Ministry of Health. These figures show RDD plots for health clinics. Plots on the left use the number of clinics as the dependent variable (intensive margin). Plots on the right use an indicator for having any positive number of clinics as the dependent variable (extensive margin).

Appendix 4 Bandwidth Sensitivity


We perform an analysis of bandwidth sensitivity and compare the results to the optimal bandwidths of [Calonico et al. \(2014b\)](#), denoted as CCT below, and [Imbens and Kalyanaraman \(2012\)](#), denoted as IK. The plots include 95% confidence intervals. We use a bandwidth of 0.5 in our main results.

Figure A8: Use of health services




Note: Authors' calculations with data from Progresa, Conapo, ENSA 2000, and the Ministry of Public Health. Plots include 95% heteroskedasticity-robust confidence intervals. CCT: [Calonico et al. \(2014a\)](#) optimal bandwidth. IK: [Imbens and Kalyanaraman \(2012\)](#) optimal bandwidth.

Figure A9: Use of health services by ages




Note: Authors' calculations with data from Progresa, Conapo, ENSA 2000, and the Ministry of Public Health. Plots include 95% heteroskedasticity-robust confidence intervals. CCT: Calonico et al. (2014a) optimal bandwidth. IK: Imbens and Kalyanaraman (2012) optimal bandwidth.

Figure A10: Use of health services with Administrative Data




Note: Authors' calculations with data from Progresa, Conapo, ENSA 2000, and the Ministry of Public Health. Plots include 95% heteroskedasticity-robust confidence intervals. CCT: Calonico et al. (2014a) optimal bandwidth. IK: Imbens and Kalyanaraman (2012) optimal bandwidth.

Figure A11: **Reproductive health**




Note: Authors' calculations with data from Progresa, Conapo, ENSA 2000, and the Ministry of Public Health. Plots include 95% heteroskedasticity-robust confidence intervals. CCT: [Calonico et al. \(2014a\)](#) optimal bandwidth. IK: [Imbens and Kalyanaraman \(2012\)](#) optimal bandwidth.

Figure A12: **Chronic disease and prevention**



Note: Authors' calculations with data from Progresa, Conapo, ENSA 2000, and the Ministry of Public Health. Plots include 95% heteroskedasticity-robust confidence intervals. CCT: [Calonico et al. \(2014a\)](#) optimal bandwidth. IK: [Imbens and Kalyanaraman \(2012\)](#) optimal bandwidth.

Figure A13: **Service characteristics**



Note: Authors' calculations with data from Progres, Conapo, ENSA 2000, and the Ministry of Public Health. Plots include 95% heteroskedasticity-robust confidence intervals. CCT: [Calonico et al. \(2014a\)](#) optimal bandwidth. IK: [Imbens and Kalyanaraman \(2012\)](#) optimal bandwidth.

Appendix 5 Additional Robustness Checks

Table A10: **Robustness of Main Results to Clustering Standard Errors by Locality**

	Variable	$\hat{\tau}_c$	Heteroskedasticity robust p-value	Locality clustered p-value	Data
(1)	ln(Total visits 2000)	0.0747	0.2223	0.2379	A
(2)	ln(Total visits 2001)	0.1293	0.0351	0.0389	A
(3)	ln(Total visits 2002)	0.1142	0.0706	0.0724	A
(4)	ln(Total visits 2003)	0.1578	0.0115	0.0116	A
(5)	Child attended by medic	0.0818	0.0338	0.0710	S
(6)	Used health services	0.0397	0.0001	0.0835	S
(7)	Self-medicated	-0.0085	0.0136	0.1860	S
(8)	Health improvement	0.0664	0.0000	0.0352	S
(9)	0–4 yo user	0.0557	0.0776	0.2503	S
(10)	5–14 yo user	0.0433	0.0163	0.1236	S
(11)	15–19 yo user	0.0084	0.7516	0.7447	S
(12)	20–49 yo user	0.0368	0.0206	0.1379	S
(13)	50–64 yo user	0.0531	0.1242	0.2389	S
(14)	65+ yo user	-0.0081	0.8527	0.8789	S
(15)	Contraceptives	0.0755	0.0155	0.0815	S
(16)	Prenatal checkups	0.5530	0.0627	0.1671	S
(17)	Has been pregnant	0.0078	0.8062	0.8150	S
(18)	Pregnant in 2000	0.0044	0.7630	0.7560	S
(19)	Diabetes diagnostic	-0.0009	0.9405	0.9445	S
(20)	HBP diagnostic	0.0318	0.0989	0.0863	S
(21)	Pap smear test	0.0076	0.5566	0.6336	S
(22)	Breast cancer test	0.0199	0.1864	0.2721	S
(23)	Waited 0–50 mins.	-0.0299	0.0478	0.3461	S
(24)	Satisfaction with waiting time	-3.0930	0.1570	0.1552	S
(25)	Complaint	0.0252	0.3671	0.3646	S
(26)	Medical explanation	-1.3520	0.4491	0.4471	S


Note: This table replicates Tables 2, 3, 4, 5, and 6 in the main text. Additional to reporting heteroskedasticity-robust p-values, we report p-values that correspond to locality-clustered standard errors. In the last column, “A” stands for administrative data, and “S” stands for survey data. Statistics are reported for 2000, except rows (2)-(4) which also report information for 2001-2003, and rows (24)-(26) which report information for 2003.

Table A11: **Robustness of Main Results to Constructing Indices from Main Variables**

	Use of health services	Reproductive health	Chronic disease and prevention		Service characteristics
			Men	Women	
$\hat{\tau}_c$	0.2946 (0.1266)	0.0868 (0.1109)	0.0154 (0.1034)	0.0769 (0.0649)	-0.2136 (0.1680)
Observations	8200	9148	10158	24186	2375
P-value	0.0200	0.4340	0.8810	0.2360	0.2030
Mean dependent variable	0.0000	0.0000	0.0000	0.0000	0.0000
Mean dep. var. left of cutoff	0.0792	-0.1730	0.1090	-0.0598	0.1280
Table	2	4	5	5	6

Note: This table shows the RDD results for indices that collapse the variables in Tables [2](#), [4](#), [5](#), and [6](#) in the main text into a single variable using principal component analysis.

Figure A14: **Robustness of Main Results to Constructing Indices from Main Variables**



Note: These figures show RDD plots for Principal Component Analysis for variables in Tables [2](#), [4](#), [5](#), and [6](#). We present binned means of the outcome variables around the program eligibility cutoff and a local polynomial of degree one.

Table A12: **Robustness of Main Results to Multiple Hypothesis Testing**

	Variable	Model p-value	Resample p-value	Romano-Wolf p-value
1	Child attended by medic	0.0338	0.0396	0.0396
2	Used health services	0.0001	0.0099	0.0099
3	Self-medicated	0.0136	0.0099	0.0297
4	Health improvement	0.0000	0.0099	0.0099
5	0-4 yo user	0.0776	0.0792	0.3270
6	5-14 yo user	0.0163	0.0198	0.1290
7	15-19 yo user	0.7520	0.7820	0.9700
8	20-49 yo user	0.0206	0.0495	0.1290
9	50-64 yo user	0.1240	0.1680	0.3760
10	65+ yo user	0.8530	0.8610	0.9700
11	Contraceptives	0.0155	0.0099	0.0396
12	Prenatal checkups	0.0627	0.0891	0.2080
13	Has been pregnant	0.8060	0.7520	0.9410
14	Pregnant in 2000	0.7630	0.7820	0.9410
15	Diabetes diagnostic	0.9410	0.9700	0.9700
16	HBP diagnostic	0.0989	0.0990	0.3270
17	Pap smear test	0.5570	0.6440	0.8420
18	Breast cancer test	0.1860	0.1190	0.5150
19	Waited 0-50 mins.	0.0478	0.0792	0.0792
20	Satisfaction with waiting time	0.1570	0.1780	0.4160
21	Complaint	0.3670	0.3760	0.6340
22	Medical explanation	0.4490	0.4750	0.6340

Note: This table shows the results of multiple hypothesis testing for the results shown in Tables 2, 3, 4, 5, and 6 in the main text. This procedures use resampling methods to control the familywise error rate (FWER), that is, the probability of rejecting at least one true null hypothesis in the family of hypotheses under test. See [Clarke et al. \(2020\)](#).

Table A13: **Robustness of Main Results to Using the Optimal Bandwidth**

	Variable	$\hat{\tau}_c$	P-value	Bandwidth	Obs.
1	ln(Total visits)	0.0417	0.5644	0.3494	8926
2	ln(Total visits)	0.0388	0.6267	0.2790	8977
3	ln(Total visits)	0.0951	0.1767	0.4074	8594
4	ln(Total visits)	0.1428	0.0412	0.4038	9444
5	Child attended by medic	0.0846	0.0620	0.3787	10460
6	Used health services	0.0388	0.0000	0.6401	174566
7	Self-medicated	-0.0119	0.0078	0.3233	174092
8	Health improvement	0.0777	0.0001	0.3571	80873
9	0-4 yo user	0.0615	0.0749	0.4316	19700
10	5-14 yo user	0.0458	0.0130	0.4816	41646
11	15-19 yo user	0.0086	0.7470	0.4991	16970
12	20-49 yo user	0.0381	0.0207	0.4722	68965
13	50-64 yo user	0.0502	0.1102	0.6059	16935
14	65+ yo user	-0.0053	0.9076	0.4676	10208
15	Contraceptives	0.0776	0.0484	0.3327	22914
16	Prenatal checkups	0.5809	0.0748	0.4259	10846
17	Has been pregnant	0.0072	0.8232	0.4908	20781
18	Pregnant in 2000	0.0042	0.8068	0.3921	24104
19	Diabetes diagnostic	-0.0006	0.9606	0.5333	40517
20	HBP diagnostic	0.0333	0.0963	0.4704	34647
21	Pap smear test	0.0093	0.4989	0.4468	52537
22	Breast cancer test	0.0239	0.1686	0.3928	27431
23	Waited 0-50 mins.	-0.0388	0.0225	0.4058	79350
24	Satisfaction with waiting time	-2.7986	0.2805	0.3413	2375
25	Complaint	0.0209	0.4843	0.4386	2375
26	Medical explanation	-1.2361	0.5074	0.4513	2375

Note: This table replicates Tables 2, 3, 4, 5, and 6 in the main text. The difference is that it considers the optimal bandwidth for each variable. The optimal bandwidth was calculated with the *rd bwselect* command, see [Calonico et al. \(2014b\)](#).

Table A14: **Robustness of Main Results to Alternative Identification of Regional Threshold Values**

	Variable	$\hat{\tau}_c$	P-value	Obs.
1	ln(Total visits)	0.0890	0.2081	8768
2	ln(Total visits)	0.1459	0.0266	8835
3	ln(Total visits)	0.1579	0.0198	8502
4	ln(Total visits)	0.1872	0.0052	9343
5	Child attended by medic	0.0302	0.4850	10356
6	Used health services	0.0224	0.0514	172777
7	Self-medicated	-0.0145	0.0003	172316
8	Health improvement	0.0569	0.0016	80112
9	0-4 yo user	0.0159	0.6634	19499
10	5-14 yo user	0.0266	0.2048	41207
11	15-19 yo user	0.0002	0.9937	16789
12	20-49 yo user	0.0228	0.2119	68320
13	50-64 yo user	0.0531	0.1875	16747
14	65+ yo user	-0.0289	0.5668	10078
15	Contraceptives	0.0907	0.0116	22695
16	Prenatal checkups	0.1052	0.7605	10746
17	Has been pregnant	0.0191	0.6002	20584
18	Pregnant in 2000	0.0049	0.7784	23859
19	Diabetes diagnostic	-0.0006	0.9658	40100
20	HBP diagnostic	0.0444	0.0484	34300
21	Pap smear test	0.0068	0.6596	52002
22	Breast cancer test	0.0345	0.0543	27153
23	Waited 0-50 mins.	-0.0163	0.3458	78611
24	Satisfaction with waiting time	-5.3897	0.0123	2354
25	Complaint	0.0192	0.5296	2354
26	Medical explanation	-3.2734	0.0501	2354

Note: This table replicates Tables 2, 3, 4, 5, and 6 in the main text. The difference is that the threshold is calculated with a quartic polynomial, as done in Card et al. (2008).